

*Dedicated to Prof. Mihai Gavrilă's 80<sup>th</sup> Anniversary*

## A NEW UNIFORM ASYMPTOTIC APPROXIMATION OF 3D SCATTERING WAVE FUNCTION FOR A CENTRAL POTENTIAL

N. GRAMA, C. GRAMA and I. ZAMFIRESCU

Horia Hulubei National Institute for Nuclear Physics and Engineering, P.O.Box MG-6, RO-077125  
Bucharest-Magurele, Romania, E-mail: grama@theory.nipne.ro

*Received December 20, 2009*

A new uniform asymptotic approximation of the 3D scattering wave function for a central potential is obtained by using an etalon function generated by the problem. The regular reduced radial wave function of zero orbital angular momentum is used as etalon function rather than a standard special function, as used in the Kravtsov-Ludwig uniform approximation. The method is applied to the scattering by a pure Coulomb potential. The obtained uniform asymptotic approximation of the 3D Coulomb scattering wave function has the same expression as the exact solution.

*Key words: 3D scattering wave function, asymptotic approximation.*

### INTRODUCTION

The knowledge in the physics of molecules, atoms and nuclei can be obtained by the study of quantum collision processes. In order to solve the scattering problem for a central potential the most frequently used method is the partial wave expansion of the solution [1, 2, 3]. However it is well known that even for problems that have been solved exactly it often happens that this solution is too complicated to be used in practical applications. Moreover, the partial wave expansion does not emphasize the functional dependence of the wave function and of the scattering amplitude on the potential parameters, which is necessary if one desires to go behind the formula and to obtain a description of what is happening during the collision. Because of this, considerable effort has been devoted to the study of approximate solution methods. The method of WKB approximation is applied in almost all branches of physics, from tides and earthquakes through optics to ionospheric theory and quantum mechanics.

The 3D WKB approximation is the main approximation used in the study of the scattering processes. This is a non-uniform asymptotic approximation, obtained by taking as a small parameter the Planck constant  $\hbar$  [4].

The advantage of the 3D WKB approximation is its high precision. The main merit of the 3D WKB approximation is that it is a "doubly asymptotic expansion" [5, 6]. This is an approximation that has asymptotic character not only for small  $\hbar$ , but also with respect to large  $|\vec{r}|$ . The remainder of this approximation is small both for small  $\hbar$  and large  $|\vec{r}|$ . The knowledge of this doubly asymptotic approximation is extremely helpful. This allows to describe the solutions in a unique fashion and to select the scattering solution that satisfies the radiation condition at infinity.

The main shortcoming of the WKB approximation is that it becomes infinite on certain surfaces, called caustics. Caustics are the higher dimensional analogue for partial differential equations of the turning points of the WKB theory for ordinary differential equations. The problem of caustic fields is topical in the scattering of light and particles (e.g. rainbow phenomena in optics, in atomic and nuclear physics), in the wave propagation (e.g. radio and microwave antenna, problems of eigenmodes and waves in cavities and waveguides), etc. Caustics as physical objects have been analyzed by Kravtsov [7].

In order to overcome the above mentioned shortcoming of the WKB approximation a different approximation that is uniformly valid in a region containing one or more caustics is necessary. The asymptotic representation of the solution is greatly simplified if one uses a uniform asymptotic approximation. The Stokes phenomenon is obviated, and the solution is represented by a single form over the entire range of variables [5, 6]. Thus a uniform asymptotic approximation provides a smooth transition across a caustic. Kravtsov [8] and Ludwig [9] derived uniform asymptotic approximation of the solution in a region containing the simplest caustics. They have represented the solution of the problem in terms of the solution of a "related" or "comparison" equation, by a method introduced and developed by Langer [10]. The method of "comparison" equation has been discussed in details by Olver [11]. Applications of the comparison equation method to the wave theory are given in [12]. In the case of the smooth convex caustic the Airy function equation has been taken as comparison equation [8, 9]. In other terms, the solution has been represented as a linear combination of Airy function and its derivative. In the vicinity of the caustics the wave function has a smooth transition between oscillatory and exponentially damped behavior, with large values on the caustic itself. The term of the solution which involves the derivative of the Airy function is relatively small near the caustics, but prominent away from caustics. This term makes the uniform expansion possible [9, 13]. Away from the caustic the solution reduces to the 3D WKB approximation.

The uniform asymptotic approximations have been used in a large variety of problems, each problem imposing the type of special functions to be employed. The generalized Bessel and Hankel functions, Airy function, Weber functions, Fresnel integral functions have been used by a series of authors [14, 15, 16, 17, 18, 19, 20, 21, 22, 23] in order to give the asymptotic solution for phenomena such as

the scattering by a general convex obstacle, creeping waves, gallery modes, edge diffraction, trapped waves, surface waves on fluid, wave propagation in a rotating ocean. More references about uniform asymptotic approximation can be found in the books [13, 24, 25, 26].

While specific problems have been solved by the uniform asymptotic approximation method, a general theory for the 3D scattering problem has not been given, in contrast to the 1D case where there is the theory for the turning points problem [5, 6]. There is no general method to determine the appropriate uniform expansion for any given problem. Catastrophe theory [13] paved the way for a general theory of uniform asymptotics based on standard integrals. However there is no general method to determine the appropriate "Ansatz" about the form of the solution *a priori*. The method of uniform asymptotic approximation has some characteristic difficulties [27]. An accurate guess ("Ansatz") for the form of the solution is necessary, otherwise singularities will occur in some undetermined coefficients. Another difficulty of the uniform asymptotic approximation is that the labor involved increases sharply as the complexity of the problem increases [27].

The success of the uniform asymptotic expansions is connected with the principle of parsimony [28]: it is important not to take too many terms in the expansion (usually not more than one term), and it is important not to include more complications in the asymptotics than are absolutely necessary. This parsimony principle makes the method of uniform asymptotic expansion easy to apply, and makes it insensitive to the fine features of the structure under consideration.

In the present work a uniform asymptotic expansion with one term of the 3D wave function describing the scattering process of a spinless particle by a central potential is obtained. In the study of the scattering process the uniform asymptotic approximation of the wave function is very important because its asymptotic at  $r \rightarrow \infty$  allows to obtain the explicit expression of the scattering amplitude, emphasizing the dependence on the potential parameters. The paper is based on a new Ansatz. Instead of using the standard special functions (Airy, Bessel, Weber, etc) as solutions of the comparison equation a function generated by the problem itself is used. The special function used in this work is the regular reduced radial wave function of zero orbital angular momentum. The uniform asymptotic approximation of the 3D scattering wave function allows finding the scattering amplitude without using the partial waves expansion. In this way the convergence difficulties related to the scattering amplitude expansion in Legendre polynomials are obviated [29]. The method is applied to a pure Coulomb potential. In [30] a uniform asymptotic approximation of the Coulomb wave function based on the Airy function was given. This approximation is valid in the region of the caustic  $\xi = 4\eta/k$ , but fails at the caustic  $\xi = 0$ , where  $\xi = r(1 - \cos\theta)$  is a parabolic coordinate. In contrast, the uniform asymptotic approximation obtained in the present paper is valid over the whole range of the variables, including the caustic  $\xi = 0$ .

The paper is organized as follows: In Section 2 a uniform asymptotic approximation for the 3D scattering wave function by a central potential is deduced by using a new Ansatz. The special function of the approximation is the regular reduced radial wave function of zero angular momentum. In Section 3 the 3D uniform asymptotic approximation of the Coulomb scattering wave function is obtained. The 3D WKB asymptotic approximation is deduced for the Coulomb potential in order to identify and analyze the caustics of this problem. This allows to choose the appropriate comparison equation in order to obtain the 3D uniform asymptotic approximation, according to the criteria to be fulfilled by the comparison equation method presented in the Appendix A. The uniform asymptotic approximation of the 3D Coulomb scattering wave function is obtained by using the new Ansatz for the form of the solution. The uniform asymptotic approximate wave function is in fact the exact Gordon solution expressed in the form obtained by Grama, Grama and Zamfirescu in a previous paper [30]. This demonstrates the accuracy of the guess (Ansatz) of the solution.

## 2. UNIFORM ASYMPTOTIC APPROXIMATION OF 3D SCATTERING WAVE FUNCTION FOR A CENTRAL POTENTIAL. NEW ANSATZ

We consider the scattering of a spinless particle by a potential  $U(\vec{r})$  that decreases at infinity. The wave function  $\Psi$  that describes the scattering process satisfies the Schrödinger equation

$$\Delta\Psi + (k^2 - V(\vec{r}))\Psi = 0 \quad (1)$$

with  $k^2 = \frac{2m}{\hbar^2}E$ ,  $V(\vec{r}) = \frac{2m}{\hbar^2}U(\vec{r})$ .  $\Psi$  is a regular function in  $\mathbf{R}^3$  and

$$\Psi = \text{plane wave} + \phi(r), \quad (2)$$

where  $\phi(r)$  satisfies the radiation condition at infinity. If  $V(\vec{r})$  is a short range potential, *i.e.*  $\lim_{r \rightarrow \infty} V(\vec{r}) \sim \mathcal{O}(r^{-\epsilon})$ ,  $\epsilon > 2$ , then the radiation condition is [31]

$$\lim_{r \rightarrow \infty} (\partial\phi/\partial r - ik\phi) = 0. \quad (3)$$

If  $V(\vec{r})$  is a long-range potential, *i.e.*  $\lim_{r \rightarrow \infty} V(\vec{r}) \sim \mathcal{O}(r^{-\epsilon})$ ,  $\epsilon < 2$ , then the radiation condition is "nonspherical" [32, 33]

$$\lim_{r \rightarrow \infty} (\nabla\phi - ik\nabla S\phi) = 0, \quad (4)$$

where  $S$  satisfies the eikonal equation

$$(\nabla S)^2 = 1 - \frac{V(\vec{r})}{k^2}. \quad (5)$$

We shall find the asymptotic solution of (1) for small Plank constant  $\hbar$ . Instead of working in (1) with small  $\hbar$  a large parameter  $\lambda$  is introduced. In other words the asymptotic solution for large  $\lambda$  of the following equation will be obtained

$$\Delta\Psi + \lambda^2(k^2 - V(\vec{r}))\Psi = 0. \quad (6)$$

In the final expression of the wave function the parameter  $\lambda$  will be put  $\lambda = 1$ . Let us consider the 3D WKB approximation of the solution for the scattering by a central potential  $V(r)$ . In order to obtain the 3D WKB approximation for  $\Psi$  we look for a solution of (6) for large  $\lambda$  of the form [4, 34]

$$\Psi = A(\vec{r})e^{i\lambda S(\vec{r})}. \quad (7)$$

Substituting (7) in (6) and by grouping the like-power terms the following equations are obtained

$$(\nabla S)^2 = k^2 - V(r) \quad \text{eikonal equation,} \quad (8)$$

$$2\nabla A \nabla S + A \nabla^2 S = 0 \quad \text{transport equation.} \quad (9)$$

By solving (8) and (9) the eikonal  $S(\vec{r})$  and the amplitude  $A(\vec{r})$  are obtained, *i.e.* the 3D WKB approximation is obtained.

Our problem is to determine a uniform asymptotic approximation of the 3D solution of the Schrödinger equation which describes the scattering by a central potential  $V(r)$ . Taking into account the characteristic difficulties of a uniform asymptotic approximation mentioned in the Introduction we introduce a new initial guess ("new Ansatz"). We start from the idea that the special function used in the form of the solution must be generated by the problem itself, rather than using standard special functions (Airy, Bessel, Weber, etc). Let us consider the partial waves expansion of the wave function  $\Psi(\vec{r})$  which describes the scattering

$$\Psi(\vec{r}) = \sum_{l=0}^{\infty} C_l \frac{R_l(r)}{r} P_l(\cos \theta), \quad (10)$$

where  $R_l(r)$  are the regular reduced radial wave functions and  $P_l(\cos \theta)$  are the Legendre polynomials. Let us take the WKB approximation of both parts of (10). Then the singularities of the WKB approximations of the regular reduced radial wave functions at the turning points and the singularities must be somehow reflected in the singularities of the 3D WKB approximation of  $\Psi(\vec{r})$  at caustics.

This leads us to the new Ansatz, namely we take as the special function which occurs in the uniform approximation of the solution  $\Psi(\vec{r})$  the regular reduced radial wave function of zero orbital angular momentum  $R_0(r)$ . In other words we take the form

$$\Psi(\vec{r}) = e^{i\lambda\omega(\vec{r})} \left\{ g(\vec{r}) R_0(\rho(\vec{r})) - \frac{i}{\lambda} h(\vec{r}) R_0'(\rho(\vec{r})) \right\}. \quad (11)$$

Here  $R_0$  is the regular solution of the second order ordinary differential equation

$$\frac{d^2 R_0}{d\rho^2} + \lambda^2 P(\rho) R_0 = 0, \quad (12)$$

where

$$P(\rho) = 1 - \frac{1}{k^2} V(\rho/k). \quad (13)$$

In order to obtain the uniform asymptotic approximation the functions  $\omega(\vec{r})$ ,  $\rho(\vec{r})$ ,  $g(\vec{r})$ , and  $h(\vec{r})$  have to be determined.

In Appendix B it is shown that the functions  $\omega(\vec{r})$ ,  $\rho(\vec{r})$ ,  $g(\vec{r})$ , and  $h(\vec{r})$  can be expressed in terms of new functions  $S_{\pm}$ ,  $\mathcal{A}_{\pm}$  in the following way

$$\omega = \frac{1}{2}(S_+ + S_-), \quad (14)$$

$$\int^{\rho} P^{1/2}(t) dt = \frac{1}{2}(S_+ - S_-), \quad (15)$$

$$g = \frac{1}{2} P^{1/4} (\mathcal{A}_+ + \mathcal{A}_-), \quad (16)$$

$$h = \frac{1}{2} P^{-1/4} (\mathcal{A}_+ - \mathcal{A}_-), \quad (17)$$

where  $S_{\pm}$ , and  $\mathcal{A}_{\pm}$  satisfy the generalized eikonal equations and generalized transport equations, respectively

$$(\nabla S_{\pm})^2 = k^2 - V(r), \quad (18)$$

$$2\nabla S_{\pm} \nabla \mathcal{A}_{\pm} + \mathcal{A}_{\pm} \Delta S_{\pm} = 0. \quad (19)$$

In order to find the uniform asymptotic approximation of the scattering solution of the Schrödinger equation (1) for a central potential  $V(r)$  using the new Ansatz the generalized eikonal and transport equations (18) and (19) have to be

solved, then their solutions  $\mathcal{S}_\pm$ ,  $\mathcal{A}_\pm$  used to determine the functions  $\omega(\vec{r})$ ,  $\rho(\vec{r})$ ,  $g(\vec{r})$ , and  $h(\vec{r})$  according to (14), (15), (16), and (17). With the functions  $\omega(\vec{r})$ ,  $\rho(\vec{r})$ ,  $g(\vec{r})$ , and  $h(\vec{r})$  the uniform asymptotic approximation of the solution of (1) is constructed in the form given by (11). The new Ansatz (11) uses the regular reduced radial wave function of zero orbital angular momentum  $R_0$  and its derivative, rather than standard special functions. In this way the pattern of the caustics of the 3D WKB approximation is the same as the pattern of the caustics of the regular radial wave functions. Indeed, the signature of the caustics of the radial wave equation (turning points and singularities) is preserved by the caustics of the 3D problem because the comparison function  $R_0(\rho(\vec{r}))$  satisfies the zero angular radial equation written in the variable  $\rho(\vec{r})$ . Moreover the new Ansatz ensures the correct behavior at infinity, *i.e.* the wave function satisfies the radiation condition at infinity. The uniform asymptotic approximation must be multiplied by a constant that is determined from the condition to have an incident wave of unit amplitude. The uniform asymptotic approach obviates the shortcomings of the partial wave expansion that consist in the impossibility to bring into prominence the dependence of the wave function on the potential parameters and the poor convergence of the expansion for large  $k$ . Instead of the series of the partial wave expansion a closed form expression for the wave function is obtained. The asymptotics at  $r \rightarrow \infty$  of the uniform asymptotic approximation of the scattering wave function leads to the explicit expression of the scattering amplitude, emphasizing the dependence on the potential parameters.

### 3. UNIFORM ASYMPTOTIC APPROXIMATION OF 3D COULOMB SCATTERING WAVE FUNCTION BASED ON A NEW ANSATZ

The 3D WKB approximation is necessary in order to identify and analyze the caustics of the problem. This allows to choose an appropriate comparison equation (see Appendix A). The analysis of the caustics characteristics shows that the new introduced Ansatz is suited for the present problem. Moreover, in order to find the uniform asymptotic approximation, the solutions of the eikonal and transport equations that occur in obtaining the 3D WKB approximation are needed. In [35] a detailed deduction of the 3D WKB asymptotic approximations for the Coulomb scattering wave function is given. The result is

$$\Psi_{in} \sim A_- e^{iS_-}, \quad \Psi_{out} \sim A_+ e^{iS_+}, \quad (20)$$

where  $A_\pm$  are the solutions of the transport equation

$$A_\pm = \frac{1 \mp w}{2w^{1/2}} \quad \text{with} \quad w = \sqrt{1 - \frac{4c}{k^2 \xi}} \quad (21)$$

and  $S_{\pm}$  are the solutions of the eikonal equation

$$S_{\pm} = \frac{k}{2} \left( \zeta \pm \int^{\xi} \sqrt{1 - \frac{4c}{k^2 u}} du \right). \quad (22)$$

For  $c = 0$  we have  $A_- = 1$ ,  $A_+ = 0$ ,  $S_- = kz$ ,  $S_+ = kr$ . It results  $\Psi_{in} = e^{ikz}$ ,  $\Psi_{out} = 0$ . For  $c \neq 0$  the function  $\Psi_{in}$  is the Coulomb distorted plane wave function and  $\Psi_{out}$  is the Coulomb distorted outgoing wave function.

Taking into account the WKB is a doubly asymptotic approximation we may take the asymptotics for  $r \rightarrow \infty$  of  $\Psi_{in}$  and  $\Psi_{out}$ . By calculating the integral in (22) and by introducing the result in the equations (20) one obtains

$$\Psi_{in} \underset{r \rightarrow \infty}{\sim} e^{i(kz + \eta \ln k \xi)}, \quad \Psi_{out} \underset{r \rightarrow \infty}{\sim} -\frac{\eta}{k(1 - \cos \theta)} \frac{e^{i(kr - \eta \ln k \xi)}}{r}. \quad (23)$$

It is to be noticed that from (23) the Rutherford scattering amplitude results up to a constant

$$a_c \sim K \frac{\eta}{2k \sin^2 \theta/2}, \quad \theta \neq 0. \quad (24)$$

The constant  $K$  in the above equation cannot be determined in the WKB

approximation. If  $\xi = \frac{4\eta}{k}$  or  $\xi = 0$ , then  $A_{\pm} \rightarrow \infty$  and the 3D WKB approximation breaks down. In other words the 3D WKB approximation has caustics at  $\xi = 0$ , and

$\xi = \frac{4\eta}{k}$ . The 3D WKB approximation of the Coulomb scattering wave function fails at the caustics, because the amplitudes  $A_{\pm}$  become infinite there. The caustic represented by the surface of the paraboloid  $\xi = \frac{4\eta}{k}$  separates the classically

forbidden region  $\left(0 < \xi < \frac{4\eta}{k}\right)$  from the classically allowed region  $\left(\xi > \frac{4\eta}{k}\right)$ . The

caustic  $\xi = 0$  is the Oz axis. The axial caustic (Oz axis) is characteristic to the solution of scattering problems that do not depend on the azimuthal angle [7, 13]. Although the 3D WKB approximation of the wave function is singular at the caustics, the actual wave function is finite there.

In order to overcome the mentioned shortcoming of the 3D WKB approximation a uniform asymptotic approximation, valid near and away from the caustics, is necessary. The uniform asymptotic approximation reduces to the 3D WKB approximation away from the caustics, and remains finite at the caustics. We are looking for a uniform asymptotic approximation for 3D Coulomb scattering wave function based on the new Ansatz of the form

$$\Psi \sim e^{i\lambda\omega(\vec{r})} \left\{ g(\vec{r}) R_0(\rho(\vec{r})) - \frac{i}{\lambda} h(\vec{r}) R_0'(\rho(\vec{r})) \right\}, \quad (25)$$

where  $R_0$  is the regular solution of the reduced radial equation for zero angular momentum

$$\frac{d^2 R_0}{d\rho^2} + \lambda^2 \left( 1 - \frac{2\eta}{\rho} \right) R_0 = 0. \quad (26)$$

In order to determine the functions  $\omega(\vec{r})$ ,  $\rho(\vec{r})$ ,  $g(\vec{r})$ , and  $h(\vec{r})$  we need the solutions  $\mathcal{S}_\pm$  and  $\mathcal{A}_\pm$  of the generalized eikonal and transport equations (18) and (19). Taking into account that equations (18) and (19) for  $\mathcal{S}_\pm$  and  $\mathcal{A}_\pm$  are identical to equations (8) and (9) for  $S_\pm$  and  $A_\pm$  it results  $\mathcal{S}_\pm = S_\pm$ ,  $\mathcal{A}_\pm = A_\pm$ . Consequently we determine  $\omega(\vec{r})$ ,  $\rho(\vec{r})$ ,  $g(\vec{r})$ , and  $h(\vec{r})$  by using  $S_\pm$  and  $A_\pm$  given by equations (22), (21). By introducing the functions  $S_\pm$  and  $A_\pm$  in the equations (14), (15), (16), (17) the functions  $\omega = \frac{k}{2}\zeta$ ,  $\rho = \frac{k}{2}\xi$ ,  $g = \frac{1}{2}$ , and  $h = -\frac{1}{2}$  obtained. In this way the uniform asymptotic approximation of the 3D Coulomb scattering wave function is obtained up to a constant factor

$$\Psi \sim \frac{1}{2} e^{\frac{1}{2}k\zeta} \left\{ R_0\left(\frac{1}{2}k\xi\right) + iR_0'\left(\frac{1}{2}k\xi\right) \right\}, \quad (27)$$

where  $R_0\left(\frac{1}{2}k\xi\right) = F_0\left(\frac{1}{2}k\xi\right)$  is the regular Coulomb wave function for zero angular momentum [37], *i.e.*  $F_0(\rho)$  is the regular solution of the differential equation

$$\frac{d^2 F_0}{d\rho^2} + \left( 1 - \frac{2\eta}{\rho} \right) F_0 = 0. \quad (28)$$

We have seen that one of the caustics is the convex surface of the paraboloid  $\xi = 4\eta/k$  having Oz as axis of symmetry and the focus in the origin. This caustic separates the classically forbidden region  $\left( 0 < \xi < \frac{4\eta}{k} \right)$  from the classically allowed region  $\left( \xi > \frac{4\eta}{k} \right)$ . The classical trajectories of the charged particles scattered by the Coulomb potential are situated in the last region and are tangent to the caustic surface [36].

We consider the uniform asymptotic approximation of the Coulomb scattering obtained above

$$\Psi \sim \frac{1}{2} e^{i\omega} \{F_0(\rho) + iF_0'(\rho)\}. \quad (29)$$

We will show that through the WKB approximation of this function we will obtain a linear combination of the 3D WKB approximations of the Coulomb distorted plane wave and Coulomb distorted outgoing wave function obtained above. The 1D WKB approximation of the Coulomb function and their derivative valid for  $\rho > 2\eta$  [37] is

$$F_0(\rho) \sim g^{-1/4} \sin(\phi + \pi/4), \quad (30)$$

$$F_0'(\rho) \sim g^{1/4} \cos(\phi + \pi/4), \quad (31)$$

where

$$\phi = \int_{2\eta}^{\rho} \sqrt{1 - \frac{2\eta}{u}} du, \quad g = 1 - \frac{2\eta}{\rho}. \quad (32)$$

By introducing (30), (31) in (29) it results

$$\Psi \sim \frac{i}{4} e^{i\omega} \left[ (g^{1/4} - g^{-1/4}) e^{i(\phi + \pi/4)} + (g^{1/4} + g^{-1/4}) e^{-i(\phi + \pi/4)} \right]. \quad (33)$$

By using  $\omega = k\zeta/2$  and  $\rho = k\xi/2$  one obtains

$$\Psi \sim \frac{i}{4} \left( -A_+ e^{iS_+} e^{i\pi/4} + A_- e^{iS_-} e^{-i\pi/4} \right). \quad (34)$$

It results that by taking the WKB approximation of the uniform asymptotic approximation for  $\xi > \frac{4\eta}{k}$  a linear combination of the 3D WKB approximations of the Coulomb distorted plane wave and Coulomb distorted outgoing wave is obtained [see (20)], in other words we see that away from caustic the uniform asymptotic approximation of 3D Coulomb scattering wave function passes into the 3D WKB approximation.

In order to have an incident Coulomb distorted plane wave of unit amplitude the uniform asymptotic approximation (29) must be multiplied by a constant. In order to determine this constant we need the asymptotic approximation for large argument of the Coulomb wave function  $F_0$  and of its derivative.

Wheeler [38] has applied the phase-amplitude method to Coulomb wave functions  $F_0(\rho)$  and  $G_0(\rho)$ , writing

$$F_0(\rho) = A_0(\rho) \sin \phi_0(\rho), \quad (35)$$

$$G_0(\rho) = A_0(\rho) \cos \phi_0(\rho). \quad (36)$$

The two terms asymptotic approximation for large  $\rho$  of  $A_0$  and  $\phi_0$  is [38]

$$A_0 = 1 + \frac{\eta}{2\rho}, \quad (37)$$

$$\phi_0 = \rho - \eta \ln 2\rho + \sigma_0, \quad (38)$$

where  $\sigma_0 = \arg \Gamma(1 + i\eta)$  (see also [37]).

The Wronskian of the approximations of  $F_0$  and  $G_0$  is unit, as it should be, since the Wronskian of the exact solutions is unit. This is an important merit of these approximations and taking the multiplicative constant in front of the uniform asymptotic approximation equal to  $2ie^{i\sigma_0}$  we find

$$\Psi \sim e^{i(kz + \eta \ln k\xi)} - \frac{\eta \Gamma(1 + i\eta)}{k \Gamma(1 - i\eta)} \frac{e^{i(kr - \eta \ln k\xi)}}{\xi} \quad (39)$$

From this relation the well-known Coulomb scattering amplitude can be deduced

$$f_c(\theta) = -\frac{\eta}{2k \sin^2 \theta/2} \frac{\Gamma(1 + i\eta)}{\Gamma(1 - i\eta)} e^{-i\eta \ln \sin^2 \theta/2}, \quad \theta \neq 0. \quad (40)$$

In this way the uniform asymptotic approximation obtained by using the new Ansatz is

$$\Psi = e^{i\sigma_0} e^{\frac{i}{2}k\xi} \left[ F_0' \left( \frac{1}{2}k\xi \right) - iF_0 \left( \frac{1}{2}k\xi \right) \right], \quad (41)$$

where  $\xi$  and  $\zeta$  are the parabolic coordinates. This asymptotic approximation is valid on the whole range of the variables, including the caustics  $\xi = 0$  and  $\xi = 4\eta/k$ . In [30] a uniform asymptotic approximation of the 3D Coulomb scattering wave function has been obtained by using an Ansatz based on the Airy function. Although this approximation is valid at the paraboloid caustic  $\xi = 4\eta/k$ , it breaks down at the caustic  $\xi = 0$  (Oz axis).

The uniform asymptotic approximation (41) coincides with the Gordon solution of the Coulomb scattering problem [36]. Indeed, in [30] we expressed the Gordon solution in terms of the Coulomb wave function  $F_0$  and its derivative by using some recurrence relations of the Whittaker functions. In other words the new Ansatz based on the regular reduced radial wave function of zero orbital angular momentum provides a uniform asymptotic approximation of the scattering wave function that in the case of Coulomb potential is the exact solution.

#### 4. CONCLUSIONS

In the present paper a uniform asymptotic approximation, based on a new Ansatz, is obtained for the scattering wave function in the case of an arbitrary central potential  $V(r)$ . The new Ansatz is introduced by using a function generated by the scattering problem, rather than by using a standard special function. This function, generated by the problem itself, is the regular reduced radial wave function of zero orbital angular momentum. By expressing the 3D uniform asymptotic approximation in terms of this function we are sure that the WKB approximation of this uniform asymptotic approximation has the same structure of caustics as the caustics of the studied 3D scattering problem. In other words, this ensures that the proper Ansatz has been chosen, as discussed in Appendix A. The uniform asymptotic approximation of the scattering wave function for an arbitrary central potential led to an explicit expression of the scattering amplitude, emphasizing the dependence on the potential parameters.

The method is applied to the scattering by a pure Coulomb potential. The 3D WKB approximation of the Coulomb scattering wave function has two caustics: a paraboloid  $\xi = 4\eta/k$  and the Oz axis ( $\xi = 0$ ), where  $\xi = r(1 - \cos\theta)$  is a parabolic coordinate. By using the new Ansatz based on the regular reduced radial wave function of zero orbital angular momentum a uniform asymptotic approximation, valid on the whole range of the variables is obtained. It obviates the shortcoming of the uniform asymptotic approximation based on the Airy function [30] that is valid on the caustic  $\xi = 4\eta/k$ , but is singular at the axial caustic Oz. The uniform asymptotic approximation of the Coulomb wave function obtained in the present paper is in fact the Gordon solution of the Coulomb scattering problem expressed in terms of the Coulomb wave function  $F_0(\rho)$  and its derivative, as it has been obtained in [30]. Being valid on the whole range of the variables, the obtained uniform asymptotic approximation allows to get the scattering amplitude by using its asymptotics at  $r \rightarrow \infty$ . By using the Coulomb wave function  $F_0(\rho)$  as solution of the comparison equation the configuration of the exceptional points (turning point and singular point), that are essential for the uniform asymptotic approximation, has been preserved.

#### 5. APPENDIX A

In the following a short description of the method of comparison equation is given. This method is used in order to obtain the uniform asymptotic expansion for large parameter of the solution of the ordinary second order linear differential equation. Let us consider the differential equation

$$\frac{d^2 y(x, \lambda)}{dx^2} + \lambda^2 p(x, \lambda) y(x, \lambda) = 0. \quad (42)$$

We are looking for a uniform asymptotic expansion of  $y(x, \lambda)$  for large  $\lambda$ . Let  $w(z, \lambda)$  be the solution of the equation (comparison equation)

$$\frac{d^2 w(z, \lambda)}{dz^2} + \lambda^2 q(z, \lambda) w(z, \lambda) = 0. \quad (43)$$

We consider the asymptotic expansion of  $y(x, \lambda)$  for large  $\lambda$

$$y(x, \lambda) = A(x, \lambda) w(z, \lambda) + \frac{1}{\lambda} B(x, \lambda) \frac{dw(z, \lambda)}{dz}, \quad (44)$$

where  $A(x, \lambda)$  and  $B(x, \lambda)$  are expanded in inverse powers of  $\lambda$ . The comparison equation (43) is chosen so that it had the same configuration of turning points and singularities as the initial equation (42). If there is a turning point of first order the comparison equation is the Airy function equation. If there are two turning points of first order or one of second order, then the comparison equation is the Weber differential equation. Lynn and Keller [39] have extended the method of comparison equation for equations having an arbitrary finite number of turning points of any order. In the papers [40, 41, 42] the authors have obtained uniform asymptotic approximations of Coulomb wave functions  $F_L(\rho)$  and  $G_L(\rho)$  for large  $\eta$  by using the comparison equation method. We restrict ourselves to  $L = 0$ . The Coulomb wave function equation for  $L = 0$  is

$$\frac{d^2 F_0}{d\rho^2} + \left(1 - \frac{2\eta}{\rho}\right) F_0 = 0. \quad (45)$$

This differential equation has two exceptional points:  $\rho = 0$  and  $\rho = 2\eta$ , that are a regular singular point, and a turning point, respectively. The above-mentioned authors have obtained two uniform asymptotic approximations of  $F_0(\rho)$  and  $G_0(\rho)$  using two comparison equations: the Airy equation and the Bessel equation. The uniform approximation using Airy functions is valid for  $\rho > 2\eta$ . The uniform asymptotic approximation using Bessel functions is valid for  $\rho < 2\eta$ . (for details see [40, 41, 42]). In order to obtain a uniform asymptotic approximation valid on the whole interval  $\rho \in [0, \infty)$  a comparison differential equation should be used which has a turning point and a regular singular point. A uniform asymptotic approximation for second order linear differential equation having a simple pole and a turning point was obtained by Dunster [43]. The turning point and the singularity may become close. The comparison equation used by Dunster was the Whittaker functions equation. In other words in the case of Coulomb wave function equation the comparison equation, that gives the uniform asymptotic

approximation on the whole interval  $[0, \infty)$  is the Coulomb wave function equation, *i.e.* the uniform asymptotic approximation is the Coulomb wave function itself.

## 6. APPENDIX B

We look for the asymptotic solution for large  $\lambda$  of the Schrödinger equation

$$\Delta\Psi + \lambda^2(k^2 - V(\vec{r})) \Psi = 0. \quad (46)$$

A new Ansatz is used, namely we take as the special function which occurs in the uniform approximation of the solution  $\Psi(\vec{r})$  the regular reduced radial wave function of zero orbital angular momentum  $R_0(r)$ . In other words we take the form (25), with  $P(\rho)$  given by (13). In order to obtain the uniform asymptotic approximation the functions  $\omega(\vec{r})$ ,  $\rho(\vec{r})$ ,  $g(\vec{r})$ , and  $h(\vec{r})$  have to be determined. By substituting (25) into the Schrödinger equation (46), and by using (26)  $R_0''(\rho)$  is eliminated. Then the coefficients that multiply  $R_0(\rho)$  and  $R_0'(\rho)$  in the Schrödinger equation are separately equated to zero. In this way two equations are obtained. By collecting in these equations the coefficients with the same power of  $\lambda$  the following system of equations for  $\omega(\vec{r})$ ,  $\rho(\vec{r})$ ,  $g(\vec{r})$ , and  $h(\vec{r})$  results

$$(\nabla\omega)^2 + P(\rho)(\nabla\rho)^2 = k^2 - V(r), \quad (47)$$

$$\nabla\omega\nabla\rho = 0, \quad (48)$$

$$2\nabla\omega\nabla g + \nabla\rho Ph + 2\nabla\rho\nabla hP + g\Delta\omega + (\nabla\rho)^2 P'h = 0, \quad (49)$$

$$2\nabla g\nabla\rho + h\Delta\omega + g\Delta\rho + 2\nabla\omega\nabla h = 0. \quad (50)$$

In order to obtain the functions  $\rho(\vec{r})$  and  $\omega(\vec{r})$  the new functions  $S_{\pm}(\vec{r})$  defined as

$$S_{\pm}(\vec{r}) = \omega(\vec{r}) \pm \int^{\rho(\vec{r})} P^{1/2}(t) dt, \quad (51)$$

are introduced. By adding/subtracting the equation (47) to/from the equation (48) multiplied by  $2P^{1/2}(\rho)$  one obtains

$$(\nabla\omega + P^{1/2}\nabla\rho)^2 = k^2 - V(r), \quad (52)$$

$$(\nabla\omega - P^{1/2}\nabla\rho)^2 = k^2 - V(r). \quad (53)$$

Taking into account the definition (51) of  $S_{\pm}$  it results that equations (52) and (53) can be written as

$$(\nabla S_{\pm})^2 = k^2 - V(r), \quad (54)$$

which represents the generalized eikonal equation. In order to obtain the functions  $g(\vec{r})$  and  $h(\vec{r})$  the new functions  $\mathcal{A}_{\pm}(\vec{r})$  defined as

$$\mathcal{A}_{\pm}(\vec{r}) = P^{-1/4}(\rho) [g(\vec{r}) \pm P^{1/2}(\rho) h(\vec{r})], \quad (55)$$

are introduced. By adding/subtracting the equation (50) multiplied by  $P^{1/4}$  to/from the equation (49) multiplied by  $P^{-1/4}$  and taking into account the definitions of  $S_{\pm}$ ,  $\mathcal{A}_{+}$ ,  $S_{-}$ , and  $\mathcal{A}_{-}$  one obtains

$$2\nabla S_{+} \nabla \mathcal{A}_{+} + \mathcal{A}_{+} \Delta S_{+} = 0, \quad (56)$$

$$2\nabla S_{-} \nabla \mathcal{A}_{-} + \mathcal{A}_{-} \Delta S_{-} = 0. \quad (57)$$

Equations (56) and (57) represent the generalized transport equation.

#### REFERENCES

1. N. F. Mott and H. S. W. Massey, *The Theory of Atomic Collisions*, 3rd ed., Clarendon Press, Oxford, 1965.
2. P. G. Burke and C. J. Joachain, *Theory of Electron-Atom Collisions. Part 1: Potential Scattering*, Plenum Press, New York, 1995.
3. R. G. Newton, *Scattering Theory of Waves and Particles*, Springer-Verlag, New York, 1982.
4. M. V. Berry and K. E. Mount, *Rep. Progr. Phys.* **35**, 315 (1972).
5. W. Wasow, *Linear Turning Point Theory*, Springer-Verlag, New-York, 198.
6. M. V. Fedoryuk, *Asymptotic Analysis*, Springer-Verlag, Berlin, 1993.
7. Yu. A. Kravtsov, *Progress in Optics*, ed. E. Wolf, Elsevier Science Publishers, vol. **26**, p. 228.
8. Yu. A. Kravtsov, *Izv. VUZ Radiofiz.* **7**, 664 (1964).
9. D. Ludwig, *Comm. Pure Appl. Math.* **19**, 215 (1966).
10. R. E. Langer, *Trans. Amer. Math. Soc.* **67**, 461 (1949).
11. F. W. J. Olver, *Unsolved Problems in the Asymptotic Estimation of Special Functions*, Academic Press, New York, 1975.
12. L. A. Weinstein, *Open Resonators and Open Waveguides*, Golem Press, Boulder, Colorado, 1969.
13. Yu. A. Kravtsov and Yu. I. Orlov, *Caustics, Catastrophes and Wave Field*, Springer-Verlag, Berlin, 1999.
14. D. Ludwig, *Comm. Pure Appl. Math.* **20**, 103 (1967).
15. R. M. Lewis, N. Bleistein and D. Ludwig, *Comm. Pure Appl. Math.* **20**, 295 (1967).
16. B. Matkowsky, *PhD Thesis*, New York University, 1966.
17. Yu. A. Kravtsov, *Izv. VUZ Radiofizika* **8**, 659 (1965).
18. R. M. Lewis, J. Boersma, *J. Math. Phys.* **10**, 2291 (1969).
19. R. Smith, *Phil. Trans. Roy. Soc.* **268**, 289 (1970).
20. E. Zauderer, *J. Math. Anal. Appl. Math.* **30**, 157 (1970).
21. M. C. Shen and J. B. Keller, *SIAM J. Appl. Math.* **28**, 857 (1975).
22. M. C. Shen, *SIAM Rev.* **17**, 38 (1975).

23. J. B. Keller and D. S. Ahluwalia, *SIAM J. Appl. Math.* **25**, 583 (1973).
24. V. F. Lazutkin, *Convex biliard and eigenfunctions of Laplace operator*, Izdatel'stvo Leningradskovo Universiteta, (in Russian), 1981.
25. D. Bouche, *Asymptotic Methods in Electromagnetics* Springer-Verlag, Berlin, 1996.
26. V. A. Borovikov and B. Y. Kinber, *Geometrical Theory of Diffraction*, The Institution of Electrical Engineers, London, 1994.
27. D. Ludwig, *SIAM Rev.* **12**, 325 (1970).
28. D. Ludwig, *SIAM J. Appl. Math.* **43**, 664 (1983).
29. H. M. Nussenzweig, *Diffraction Effects in Semiclassical Scattering*, Cambridge University Press, 1992.
30. C. Grama, N. Grama, and I. Zamfirescu, *Europhys. Lett.* **59**, 166 (2002).
31. A. Sommerfeld, *Partial Differential Equations in Physics*, Academic Press, New York, 1949.
32. Y. Saito, *Pacific Journal of Mathematics* **126**, 331 (1987).
33. B. Barles, *Commun. in Partial differential equations* **12**, 263 (1987).
34. Yu. A. Kravtsov and Yu. I. Orlov, *Geometrical Optics of Inhomogeneous Media*, Springer-Verlag, Berlin (1990).
35. N. Grama, C. Grama, and I. Zamfirescu, *J. Phys. A: Math. Theor.* **42**, 465306 (2009).
36. W. Gordon *Z. Phys.* **48**, 180 (1928).
37. M. H. Hull and G. Breit, *Coulomb Wave Functions Encyclopedia of Physics* **41/1**, ed. Flüge F, Springer-Verlag, Berlin, 1959, Chapt. 2.
38. J. A. Wheeler, *Phys. Rev.* **52**, 1123 (1937).
39. R. Y. S. Lynn and J. B. Keller, *Comm. Pure Appl. Math.* **23**, 379 (1970).
40. A. Erdélyi, M. Kennedy, J. L. McGregor, and C. A. Swanson, *Technical Report Ref. NR 043-121 4*, California Institute of Technology, 1955.
41. A. Erdélyi and C. A. Swanson C, *Technical Report Ref. NR 043-121*, 1955.
42. A. Erdélyi and C. A. Swanson *Mem. Am. Math. Soc.* **25**, 1 (1957).
43. T. M. Dunster, *SIAM J. Appl. Math.* **25**, 322 (1994).